Final Technical Report

DTIC FILE COPY

DYNAMICS OF OCEANIC MOTIONS

AD-A231 136
DIC Electe
JAN 17 1991

ONR Contract N00014-84-C-0461

1 March, 1984 — 31 December, 1988

by

Allan R. Robinson

Principal Investigator

Gordon McKay Professor of Geophysical Fluid Dynamics
Division of Applied Sciences, Harvard University

This project researched the dynamics of oceanic motions: aspects of the theory and modeling of fundamental dynamical and energetic processes in the sea and their interactions, and the relationship of theory and modeling to the interpretation, analysis and design of observational data and experiments. Research was directed towards the dynamics, simulation and forecasting of the low frequency variability of ocean currents (mid-ocean eddies and intense current systems), and in the physics and dynamics of the mid-latitude general circulation. It included near surface layer/deep current interactions, and the interactions of deep ocean currents with coastal and topographic features. Our modeling research was directed toward studies of the local dynamics of (partially) open regions of the ocean (i.e., arbitrary regions with flow across their boundaries), and the relationship of the regional dynamics to the larger scale dynamics and circulation in which it is embedded. The approach was appropriate for the combined use of intensive in situ data sets in combination with extensive remotely sensed and extensive sparse in situ data sets.

Developments and applications of an embedded dynamical model hierarchy, with regional to basic scales were carried out. The hierarchy is a set of compatible and intercomparable dynamical models with differing physics allowing for efficiency in computations and physical analyses. Dynamical process studies carried out via detailed vorticity and energy analyses required optimal field estimates with dynamically consistent fields obtained via data assimilation. Because data assimilation was in its infancy in oceanography, methodological studies and dynamical process studies were carried out and interpreted together. Melded data field estimates produced via dynamical interpolation and adjustment we refer to as systematic estimates, where the components of the system are the observational network, the dynamical and statistical models, and the analysis and assimilation schemes. Substantial progress was made in developing components of the Harvard (data assimilative) dynamical model hierarchy which achieved the physical options of: quasigeostrophic (QG) mesoscale dynamics coupled with surface boundary layer (SBL) dynamics, in open or partially closed domains of arbitrary coastal shapes; or primitive equation (PE) dynamics with hybrid coordinates and arbitrarily high topography. A series of process, hindcasting, and predictability studies were: concluded for the POLYMODE Synoptic Dynamics Experiment in the Western North Atlantic; carried out for the California Current system in the OPTOMA program (joint Harvard/Naval Postgraduate School); carried out in the Gulf-Stream Meander & Ring (GSM&R) region associated with GULFCAST and SYNOP;

and initiated in the Eastern Mediterranean Sea (POEM).
This report consists of a brief summary of the research usuals and a listing of the The project produced 37 refereed publications and 17 additional technical reports which were p are indexed on attached lists. The research summary will be based primarily on the publication list since the reports are for the most part preliminary or extended versions of the publications. An exception is Report # 5 which is the Proceedings of the second Ocean Prediction Workshop (OPW-86) which I co-chaired and co-edited.

Highlight contributions include: the construction and operation in real time for 2 years of a Gulf-Stream meander and ring nowcast and forecast system (Publication # 27), the clucidation of detailed mechanisms of warm and cold core ring formations (15), the assimilation

> Statement "A" per telecon Dr. David Evans Office of Naval Research/code 1122MI VHG

Di t

Faeroe front (25), the direct coupling of real-ocean-data-based Harvard dynamical model output to an acoustic propagation model (18), the development of a primitive equation open ocean modeling capability (28), the publication of a comprehensive overview of the field of geophysical fluid dynamics (24), and an extensive summary of the Harvard quasi-geostrophic methodology (13). Two additional review and overview articles dealt with the relationship of data assimilation and dynamics (6) and the ocean prediction problem (12). There were four Ph.D. theses published (1, 14, 16, 21).

Model developments included: the theoretical basis (22) and computational implementation (34) of arbitrary coastal configurations for partially open QG regions (21); the primitive equation open ocean model (16); the 3-d surface boundary layer model coupling to the QG model (14); publication of the Harvard multivariate objective analysis models (7) and the development of an objective "contour" analysis scheme which preserves fronts and gradients (32); and a method for vertical gradient estimation (19). Data analyses in the California Current compared subsurface features with surface IR patterns (2) and determined the vertical modal structure of the mesoscale variability (3). Methodological research related to the quantification of the oceanographic usefulness of satellite altimetric data: demonstrated that assimilation into a dynamical model could achieve accurate deep fields via a simulation in the POLYMODE region (9); analyzed the synoptic signal in the historical GEOS-3 and Seasat data sets (17); and constructed a 'synthetic geoid' for absolute sea surface topographic calculations from GEOSAT by combining the mean of the GULFCAST daily synoptic estimates with the altimetric mean (23). Fundamental dynamical bases were created for the study of energy and vorticity processes ocuring in data dynamically filtered via quasigeostrophic (EVA) (5) or primitive equation (PRE-EVA) (29) dynamics. In the POLYMODE region, the potential vorticity balance was estimated (20) and an EVA analysis of the most energetic events indicated intermittent 'bursts' of baroclinic instability (10) which was modified only slightly via comparison with a PRE-EVA analysis (29). A PE-QG dynamical comparison of warm and cold ring formation

events showed basically similar processes, but identified quantitative differences due to PE asymmetries (36). This study also importantly established the validity of QG dynamics in the Gulf-Stream region for periods of 2-3 weeks even over high topography because of geostrophic contour constraints. An idealized study induced ring formation by seeding point vortices (30). Forecasting research included: a sensitivity study with OPTOMA data (11); the determination of efficient forecast schemes and their relationships to dynamical events with POLYMODE data (37); a predictability study for the NW Atlantic with POLYMODE data (8); and the establishment of the capability of accurate Gulf Stream front forecasts with adequate data for assimilation (26). Research involving the coupling of acoustic propagation models to dynamical model outputs and the propagation through realistic environmental nowcasts and forecasts resulted in publications on: interfacing (31), on the sensitivities of propagation to the ring-stream environment (33); and first results on three dimensional propagation and combined frontal/topographic effects (35). A study of the effect of the pole tide on the earth's rotation was also completed (4).

Published

- 1. Pinardi, N. (1985) Quasigeostrophic Energetics and Oceanic Mesoscale Dynamics, Harvard University, Cambridge, MA (Ph.D. Thesis).
- 2. Rienecker, M.M., C.N.K. Mooers, D.E. Hagan and A.R. Robinson (1985) A Cool Anomaly off Northern California: An Investigation Using IR Imagery and In Situ Data, Journal of Geophysical Research, 90, (C3), 4807-4818.
- 3. Smith, J.A., C.N.K. Mooers and A.R. Robinson (1985) Estimation of Baroclinic Quasi-Geostrophic Modal Amplitudes from XBT/CTD Survey Data, Journal of Atmospheric and Oceanic Technology 2(4), 491-507.
- 4. Carton, J.A. and J.M. Wahr (1986) Modelling the Pole Tide and its Effect on the Earth's Rotation, Geophysical Journal of the R.A.S. 84, 121-138.
- 5. Pinardi, N. and A.R. Robinson (1986) Quasigeostrophic Energetics of Open Ocean Regions, Dynamics of Atmospheres and Oceans 10(3), 185-221.
- 6. Robinson, A.R. (1986) Data Assimilation, Mesoscale Dynamics and Dynamical Forecasting, in *Advanced Physical Oceanographic Numerical Modelling*, 465-483 (J.J. O'Brien, ed.) R. Reidel Publishing Co., Dordrecht, The Netherlands.
- 7. Carter, E.F. and A.R. Robinson (1987) Analysis Models for the Estimation of Oceanic Fields, Journal of Atmospheric and Oceanic Technology 4, 49-74.
- 8. Carton, J.A. (1987) How Predictable are the Geostrophic Currents in the Recirculation Zone? Journal of Physical Oceanography 17(6), 751-762.
- 9. DeMey, P. and A.R. Robinson (1987) Assimilation of Altimeter Eddy Fields in a Limited Area Quasi-Geostrophic Model, Journal of Physical Oceanography 17(12), 2280-2293.
- 10. Pinardi, N. and A.R. Robinson (1987) Dynamics of Deep Thermocline Jets in the POLYMODE Region, Journal of Physical Oceanography 17, 1163-1188.
- 11. Rienecker, M.M., C.N.K. Mooers and A.R. Robinson (1987) Dynamical Interpolation and Forecast Experiments of the Evolution of Mesoscale Features off Northern California, Journal of Physical Oceanography 17(18), 1190-1213.
- 12. Robinson, A.R. (1987) Predicting Open Ocean Currents, Fronts and Eddies, Three Dimensional Ocean Models of Marine and Estuarine Dynamics 89-12 (J.C.J. Nihoul and B.M. Jamart, eds.), Elsevier Scientific Publications, Amsterdam.

- 13. Robinson, A.R. and L.J. Walstad (1987) The Harvard Open Ocean Model: Calibration and Application to Dynamical Process Forecasting and Data Assimilation Studies, Applied Numerical Mathematics 3(1-2), 89-131.
- 14. Walstad, L.J., (1987) Modelling and Forecasting of Deep Ocean and Near Surface Mesoscale Eddies: Hindcasting and Forecasting with, and Coupling a Surface Boundary Layer Model to the Harvard Quasigeostrophic Model, Harvard University, Cambridge, MA (Ph.D. thesis).
- 15. Robinson, A.R., M.A. Spall and N. Pinardi (1988) Gulf Stream Simulations and the Dynamics of Ring and Meander Processes, *Journal of Physical Oceanography* 18(12), 1811-1853.
- 16. Spall, M.A. (1988) Regional Ocean Modelling: Primitive Equation and Quasi-Geostrophic Studies, Harvard University, Cambridge, MA (Ph.D. thesis).
- 17. DeMey, P. and Y. Menard (1989) Synoptic Analysis and Dynamical Adjustment of GEOS-3 and Seasat Altimeter Eddy Fields in the Northwest Atlantic, Journal of Geophysical Research 94(C5), 6221-6230.
- 18. Lee, D., G. Botseas, W.L. Siegmann and A.R. Robinson (1989) Numerical Computations of Acoustic Propagation through Three-Dimensional Ocean and Eddies, in *Numerical and Applied Mathematics* 317-321 (W.F. Ames, ed.), J.C. Baltzer Scientific Publishing Company, Basel.
- 19. Mariano, A.J. (1989) Estimation of Mesoscale Vertical Derivatives of Potential Temperature and Density from Hydrographic Data, Journal of Atmospheric and Oceanic Technology 6(6), 1013-1023.
- 20. Mariano, A.J. and H.T. Rossby (1989) Lagrangian Potential Vorticity Balance during POLYMODE, Journal of Physical Oceanography 19(7), 927-939.
- 21. Milliff, R.M. (1989) Quasigeostrophic Ocean Flows in Coastal Domains (Ph.D. thesis), UCSB, Santa Barbara, CA, prepared at Harvard University.
- 22. Pinardi, N. and R.F. Milliff (1989) A Note on Consistent Quasigeostrophic Boundary Conditions in Partially Open, Simply and Multiple Connected Domains, Dynamics of Atmospheres and Oceans 14(1-2), 65-77.
- 23. Porter, D.L., A.R. Robinson, S.M. Glenn and E.B. Dobson (1989) The Synthetic Geoid and the Estimation of Mesoscale Absolute Topography from Altimeter Data, Johns Hopkins APL Digest 10(4), 369-379.
- 24. Robinson, A.R. (1989) Progress in Geophysical Fluid Dynamics, Earth-Science Reviews 26, 191-219.

- 25. Robinson, A.R., L.J. Walstad, J. Calman, E.B. Dobson, D.W. Denbo, S.M. Glenn, D.L Porter and J. Goldhirsh (1989) Frontal Signals East of Iceland from the GEOSAT Altimeter, Geophysical Research Letters 16(1), 77-80.
- 26. Robinson, A.R., M.A. Spall, L.J. Walstad and W.G. Leslie (1989) Data Assimilation and Dynamical Interpolation in GULFCASTING Experiments, Dynamics of Atmospheres and Oceans 13(3-4), 301-316.
- 27. Robinson, A.R., S.M. Glenn, M.A. Spall, L.J. Walstad, G.M. Gardner and W.G. Leslie (1989) Forecasting Gulf Stream Meanders and Rings, EOS, The Oceanography Report 10(45), 1464-1473.
- 28. Spall, M.A. and A.R. Robinson (1989) A New Open Ocean Hybrid Coordinate Primitive Equation Model, Mathematics and Computers in Simulation 31, 241-269.
- 29. Spall, M.A. (1989) Regional Primitive Equation Modelling and Analysis of the POLY-MODE Data Set, Dynamics of Atmospheres and Oceans 14(1-2), 125-174.
- 30. Suri, A.K. and N. Pinardi (1989) Point Vortices Embedded in Thin Jet Stream: A Model for Transition and Ring Formation Processes in the Gulf Stream, *Proceedings* of the Seventh Symposium on Turbulent Shear Flows, Stanford University.
- 31. Glenn, S.M. and A.R. Robinson (1990) Nowcasting and Forecasting of Oceanic Dynamic and Acoustic Fields II, Computational Acoustics II, 117-128 (D. Lee, A. Cakmak, R. Vichnevetsky, eds.), Elsevier Scientific Publishers, Amsterdam.
- 32. Mariano, A.J. (1990) Contour Analysis: A New Approach for Melding Geophysical Fields, Journal of Atmosphere and Oceanic Technology 7(2), 285-295.
- 33. Mellberg, L.E., A.R. Robinson and G. Botseas (1990) Modelled Time Variability of Acoustic Propagation through a Gulf Stream Meander, Journal of Acoustical Society of America 87(3), 1044-1054.
- 34. Milliff, R.M. (1990) A Modified Capacitance matrix Method to Implement Coastal Boundaries in the Harvard Open Ocean Model, Mathematics and Computers in Simulation 31(6), 541-564.
- 35. Siegmann, W.L., M.L. Jacobson, D. Lee, G. Botseas, A.R. Robinson and S.M. Glenn (1990) Interfacing Mesoscale Ocean Prediction and Parabolic Acoustic Propagation Models, Computational Acoustics, Volume II 165-168 (D. Lee, A. Cakmak and R. Vichnevetsky, eds.), Elsevier Scientific Publishers, Amsterdam.
- 36. Spall, M.A. and A.R. Robinson (1990) Regional Primitive Equation and Quasigeostrophic Studies of the Gulf Stream Meander and Ring Formation, Journal of Physical Oceanography 20(7), 985-1016.

37. Walstad, I.J. and A.R. Robinson (1990) Hindcasting and Forecasting of the POLY-MODE Data Set with the Harvard Open Ocean Model, Journal of Physical Oceanography 20(11), 1682-1702.

Reports

- 1. DeMey, P. and A.R. Robinson (1984) Simulation and Assimilation of Satellites Altimeter Data at the Oceanic Mesoscale, in Frontiers of Remote Sensing of the Troposphere from Air and Space Platforms, Proceedings of the URSI Commission F. Symposium and Workshop, Israel, May 14-25.
- 2. Pinardi, N. (1985) Quasigeostrophic Energetics and Ocean Mesoscale Dynamics, Harvard Open Ocean Model Reports 20, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA (Ph.D. thesis).
- 3. Carton, J.A. and D.J. McGillicuddy (1985) Comprehensive Maps of POLYMODE Streamfunction, Harvard Open Ocean Model Reports 21, Reports in Meteorology and Oceanography, Harvard University.
- 4. DeMey, P., J.A. Carton, Y. Menard, and D.J. McGillicuddy (1986) Comparison of Satellite Altimetry with POLYMODE Observations, Ocean Modelling 67, 1-3.
- Mooers, C.N.K., A.R. Robinson, J.D. Thompson (editors) (1987) Ocean Prediction Workshop 1986: A Status and Prospectus Report on the Scientific Basis and the Navy's Needs, Proceedings of the Ocean Prediction Workshop, Phase 1 Cambridge, MA, April 1986, Phase II Long Beach, MI, November 1986.
- 6. Robinson, A.R., M.A. Spall, W.G. Leslie, L.J. Walstad and D.J. McGillicuddy (1987) Gulfcasting: Dynamical Forecast Experiments for Gulf Stream Rings and Meanders November 1985-June 1986, Harvard Open Ocean Model Reports 22, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA.
- 7. McGillicuddy, D.J. (1987) Quasigeostrophic Modelling of Isolated Vortices, Harvard Open Ocean Model Reports 27, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA (B.A. Thesis).
- 8. Walstad, L.J. (1987) Modelling and Forecasting Deep Ocean and Near Surface Mesoscale Eddies, Hindcasting and Forecasting with, and Coupling a Surface Boundary Layer Model to, the Harvard Quasigeostrophic Model, Harvard Open Ocean Model Reports 23, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA (Ph.D. Thesis).
- 9. Glenn, S.M., A.R. Robinson and L.J. Walstad (1987) Recent Results from the Harvard Gulf Stream Forecast Program, Oceanographic Monthly Summary 7(3), 12-13.
- 10. Mariano, A.J. (1988) Space-Time Interpolation of Gulf Stream North Wall Positions, Harvard Open Ocean Model Reports 29, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA.

- 11. Spall, M.A. (1988) Regional Ocean Modelling: Primitive Equation and Quasigeostrophic Studies, Harvard Open Ocean Model Reports 28, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA (Ph.D. thesis).
- 12. McGillicuddy, D.J. (1989) One-Dimensional Numerical Simulation of New Primary Production: Lagrangian and Eulerian Formulations, Harvard Open Ocean Model Reports 36, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA.
- 13. McGillicuddy, D.J. (1989) Real Time Nowcasting and Forecasting of the Mesoscale Eddy Field in the Vicinity of the OFP Site on the Cruise CI8809, Harrard Open Ocean Model Reports 37, Reports in Methorology and Oceanography, Harvard University, Cambridge, MA.
- 14. Milliff, R.F. (1989) Quasigeostrophic Ocean Flow in Coastal Domains, Harvard Open Ocean Model Reports 34, Reports in Meteorology and Oceanography, Harvard University, Cambridge, MA (Ph.D. thesis).
- 15. Robinson, A.R. and E.B. Dobson (1989) Synoptic Variability of Frontal Meandering and Eddies from GEOSAT Altimetric Data East of Iceland, Johns Hopkins University/Applied Physics Laboratory (JHU/APL, SIR 894-035).
- Robinson, A.R. and S.M. Glenn (1989) HARVARD UNIVERSITY GULFCAST: Gulfcasts/Nogufs Model Validation Results and Appendix, Harvard University, Cambridge, MA.
- 17. Porter, D.L., S.M. Glenn and A.R. Robinson (1990) GEOSAT Observations and Validation of the Total Dynamic Topography of the Gulf Stream, Proceedings of a IAPSO/SCOR Workshop, Venice, Italy: Oceans from Space.